

# Associativity and non-associativity of some hypergraph products

Richard H. Hammack, Marc Hellmuth, Lydia Ostermeier and Peter F. Stadler

**Abstract.** Several variants of hypergraph products have been introduced as generalizations of the strong and direct products of graphs. Here we show that only some of them are associative. In addition to the Cartesian product, these are the minimal rank preserving direct product, and the normal product. Counter-examples are given for the strong product as well as the non-rank-preserving and the maximal rank preserving direct product.

## 1. Introduction

Hypergraphs are natural generalizations of undirected graphs in which “edges” may consist of more than two vertices. Products of hypergraphs have been well-investigated since the 1960s, see e.g. [1, 2, 3, 4, 6, 7, 9, 10, 12, 13, 14, 15, 18].

The article [8] surveyed the literature on hypergraph products. In addition to well-known constructions such as the Cartesian product and the square product, it also considered various generalizations of graph products that had rarely been studied, if at all. In particular, it considered several variants of hypergraph products that generalize the direct and strong product of *graphs*, namely the direct products,  $\widehat{\times}$  and  $\widetilde{\times}$ , and the strong product  $\boxtimes$ . In addition, it treated the normal product  $\boxdot$ , a generalization of the strong graph product, and the direct product  $\overline{\times}$ , which were introduced by Sonntag in the 1990’s [16, 17].

Associativity is an important property of product operators. It is e.g. implicitly assumed in the standard definition of prime factors and thus for decompositions of a given hypergraph into prime factors w.r.t. a given product [5, 11].

The survey [8] mistakenly stated that the direct products  $\widehat{\times}$  and  $\widetilde{\times}$  and the strong product  $\boxtimes$  are associative. Here we give a simple counterexample for these cases and prove associativity of the remaining hypergraph products. This contribution is an addendum to the results discussed in [8].

## 2. Preliminaries

We start our brief discussion with the formal definition of hypergraphs, and the hypergraph products in question.

A (*finite*) *hypergraph*  $H = (V, E)$  consists of a (finite) vertex set  $V$  and a collection  $E$  of non-empty subsets of  $V$ . The *rank* of a hypergraph  $H = (V, E)$  is  $r(H) = \max_{e \in E} |e|$ . A *homomorphism* from  $H_1 = (V_1, E_1)$  into  $H_2 = (V_2, E_2)$  is a map  $\phi : V_1 \rightarrow V_2$  for which  $\phi(e)$  is an edge in  $H_2$  whenever  $e$  is an edge in  $H_1$ . A bijective homomorphism  $\phi$  whose inverse is also a homomorphism is called an *isomorphism*. A hypergraph is *simple* if no edge is contained in any other edge and each edge contains two or more vertices.

In what follows, we consider six hypergraph products  $\square, \widetilde{\times}, \widehat{\times}, \widetilde{\times}, \widetilde{\boxtimes},$  and  $\widehat{\boxtimes}$ . For each of these, the vertex set of the product is the Cartesian product of the vertex sets of its factors. To be more precise, given two hypergraphs  $H_1 = (V_1, E_1)$  and  $H_2 = (V_2, E_2)$  and some product  $\otimes \in \{\square, \widetilde{\times}, \widehat{\times}, \widetilde{\times}, \widetilde{\boxtimes}, \widehat{\boxtimes}\}$ , then  $V(H_1 \otimes H_2) = V(H_1) \times V(H_2)$ . The edge sets of the various products are defined as follows.

**Cartesian Product  $\square$ :**

This is an immediate generalization of the standard Cartesian product of graphs. Its edges are

$$E(H_1 \square H_2) = \{\{x\} \times f \mid x \in V(H_1), f \in E(H_2)\} \cup \{e \times \{y\} \mid e \in E(H_1), y \in V(H_2)\}.$$

There are several ways to generalize the direct product of graphs to a product of hypergraphs. Because we want such products to coincide with the usual direct product when the factors have rank 2 (and are therefore graphs) it is necessary to impose some *rank restricting* conditions on the edges. This can be accomplished in different ways and leads to different variants of the direct and strong graph products, respectively.

**Minimal Rank Preserving Direct Product  $\widetilde{\times}$ :**

Given  $e_1 \in E_1$  and  $e_2 \in E_2$ , let  $r_{e_1, e_2}^- = \min\{|e_1|, |e_2|\}$ . The edge set of this product is defined as

$$E(H_1 \widetilde{\times} H_2) := \left\{ e \in \binom{e_1 \times e_2}{r_{e_1, e_2}^-} \mid e_i \in E_i \text{ and } |p_i(e)| = r_{e_1, e_2}^-, i = 1, 2 \right\}.$$

The edges are thus the subsets  $e \subseteq e_1 \times e_2$  (with  $e_i \in E_i$ ) for which both projections  $p_i : e \rightarrow e_i$  are injective and at least one is surjective.

**Maximal Rank Preserving Direct Product  $\widehat{\times}$ :**

Given  $e_1 \in E_1$  and  $e_2 \in E_2$ , let  $r_{e_1, e_2}^+ = \max\{|e_1|, |e_2|\}$ . The edge set of this product is defined as

$$E(H_1 \widehat{\times} H_2) := \left\{ e \in \binom{e_1 \times e_2}{r_{e_1, e_2}^+} \mid e_i \in E_i \text{ and } p_i(e) = e_i, i = 1, 2 \right\}.$$

The edges are thus the subsets  $e \subseteq e_1 \times e_2$  (with  $e_i \in E_i$ ) for which both projections  $p_i : e \rightarrow e_i$  are surjective and at least one is injective.

**Non-rank-preserving Direct Product  $\widetilde{\times}$ :**

$$E(H_1 \widetilde{\times} H_2) := \{\{(x, y)\} \cup ((e \setminus \{x\}) \times (f \setminus \{y\})) \mid x \in e \in E_1; y \in f \in E_2\}.$$

The strong product of graphs is defined as  $E(G_1 \boxtimes G_2) = E(G_1 \square G_2) \cup E(G_1 \times G_2)$ . This leads to the following generalizations to hypergraphs.

**Normal Product  $\widetilde{\boxtimes}$ :**

$$E(H_1 \widetilde{\boxtimes} H_2) = E(H_1 \square H_2) \cup E(H_1 \widetilde{\times} H_2).$$

**Strong Product  $\widehat{\boxtimes}$ :**

$$E(H_1 \widehat{\boxtimes} H_2) = E(H_1 \square H_2) \cup E(H_1 \widehat{\times} H_2).$$

### 3. Associativity and Non-associativity of Hypergraph Products

It is well known that the Cartesian product is associative [9]. In contrast, we will show below that none of the products  $\widehat{\times}, \widetilde{\times}$  and  $\widehat{\boxtimes}$  is associative. Our counterexamples require the following lemma.

**Lemma 1.** *If  $G$  and  $H$  are simple hypergraphs with  $r(G) = 2$  and  $r(H) \leq 3$ , then  $G \widehat{\times} H = G \widetilde{\times} H$ .*

*Proof.* By definition,  $V(G \widehat{\times} H) = V(G \widetilde{\times} H)$ . We need to show that  $E(G \widehat{\times} H) = E(G \widetilde{\times} H)$ . Given  $* \in \{\sim, \frown\}$  and edges  $e_1, e_2$ , let  $e_1 \overset{*}{\times} e_2$  denote the set  $E((e_1, \{e_1\}) \overset{*}{\times} (e_2, \{e_2\}))$ . Then

$$E(G \overset{*}{\times} H) = \bigcup_{e_1 \in E(G), e_2 \in E(H)} (e_1 \overset{*}{\times} e_2).$$

It suffices to show that  $e_1 \widehat{\times} e_2 = e_1 \widetilde{\times} e_2$  holds for all  $e_1 \in E(G)$  and  $e_2 \in E(H)$ . Therefore, let  $e_1 \in E(G)$  and  $e_2 \in E(H)$  and assume first  $|e_2| = 2$ . Say  $e_1 = \{x_1, y_1\}$  and  $e_2 = \{x_2, y_2\}$ . Then

$$e_1 \widehat{\times} e_2 = \{\{(x_1, x_2), (y_1, y_2)\}, \{(x_1, y_2), (y_1, x_2)\}\} = e_1 \widetilde{\times} e_2.$$

Now suppose  $|e_2| = 3$ , say  $e_2 = \{x_2, y_2, z_2\}$ . Then

$$\begin{aligned} e_1 \widehat{\times} e_2 = & \{\{(x_1, x_2), (x_1, y_2), (y_1, z_2)\}, \{(x_1, x_2), (y_1, y_2), (y_1, z_2)\}, \\ & \{(x_1, x_2), (y_1, y_2), (x_1, z_2)\}, \{(y_1, x_2), (x_1, y_2), (x_1, z_2)\}, \\ & \{(y_1, x_2), (x_1, y_2), (y_1, z_2)\}, \{(y_1, x_2), (y_1, y_2), (x_1, z_2)\}\} = e_1 \widetilde{\times} e_2. \end{aligned}$$

Thus the assertion follows.  $\square$

Now we present a counterexample showing that none of the products  $\widehat{\times}$ ,  $\widetilde{\times}$  and  $\boxtimes$  is associative.

**Counterexample.** Consider the two hypergraphs  $G = (\{a, b\}, \{\{a, b\}\})$  and  $H = (\{x, y, z\}, \{\{x, y, z\}\})$ . For  $\otimes \in \{\widehat{\times}, \widetilde{\times}, \boxtimes\}$ , we claim  $G \otimes (G \otimes H) \not\cong (G \otimes G) \otimes H$ . Put  $e = \{(a, (a, x)), (a, (b, y)), (b, (b, z))\}$ . Note that  $e$  is an edge of  $G \widehat{\times} (G \widehat{\times} H)$ , and hence also of  $G \boxtimes (G \boxtimes H)$ . However, the set  $\{((a, a), x), ((a, b), y), ((b, b), z)\}$  is not an edge in  $(G \boxtimes G) \boxtimes H$ , thus also not in  $(G \widehat{\times} G) \widehat{\times} H$ , because  $\{(a, a), (a, b), (b, b)\}$  is neither an edge in  $G \boxtimes G$  nor in  $G \widehat{\times} G$ . Thus the map  $(g, (g', h)) \mapsto ((g, g'), h)$  is not an isomorphism  $G \widehat{\times} (G \widehat{\times} H) \rightarrow (G \widehat{\times} G) \widehat{\times} H$ , nor is it an isomorphism  $G \boxtimes (G \boxtimes H) \rightarrow (G \boxtimes G) \boxtimes H$ . Moreover, the following argument shows there is no isomorphism at all.

It is shown in [7] that the number of edges in  $H_1 \widehat{\times} H_2$  is

$$|E(H_1 \widehat{\times} H_2)| = \sum_{e_1 \in E_1, e_2 \in E_2} (\min\{|e_1|, |e_2|\})! S_{\max\{|e_1|, |e_2|\}, \min\{|e_1|, |e_2|\}},$$

where  $S_{n,k} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$  is a Stirling number of the second kind. Furthermore,

$$\begin{aligned} |E(H_1 \boxtimes H_2)| &= |E(H_1 \widehat{\times} H_2)| + |E(H_1 \square H_2)| \\ &= |E(H_1 \widehat{\times} H_2)| + |V(H_1)| |E(H_2)| + |E(H_1)| |V(H_2)|. \end{aligned}$$

Using this, we see that  $|E(G \widehat{\times} (G \widehat{\times} H))| = 36 \neq 12 = |E((G \widehat{\times} G) \widehat{\times} H)|$  and  $|E(G \boxtimes (G \boxtimes H))| = 82 \neq 58 = |E((G \boxtimes G) \boxtimes H)|$ . Thus  $G \widehat{\times} (G \widehat{\times} H) \not\cong (G \widehat{\times} G) \widehat{\times} H$  and  $G \boxtimes (G \boxtimes H) \not\cong (G \boxtimes G) \boxtimes H$ .

Moreover, Lemma 1 implies  $G \widetilde{\times} (G \widetilde{\times} H) = G \widehat{\times} (G \widehat{\times} H) \neq (G \widehat{\times} G) \widehat{\times} H = (G \widetilde{\times} G) \widetilde{\times} H$ .

The remainder of this contribution proves that the direct product  $\widetilde{\times}$  and the normal product  $\boxtimes$  are associative. To our knowledge, these results have not yet appeared in the literature.

**Proposition 2.** *The direct product  $\widetilde{\times}$  is associative.*

*Proof.* Let  $H_1 = (V_1, E_1)$ ,  $H_2 = (V_2, E_2)$ , and  $H_3 = (V_3, E_3)$  be hypergraphs and consider the map  $\psi : V(H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)) \rightarrow V((H_1 \widetilde{\times} H_2) \widetilde{\times} H_3)$  defined as  $(x, (y, z)) \mapsto ((x, y), z)$ . We will show that  $\psi$  is an isomorphism. Clearly  $\psi$  is bijective. Hence it remains to show the isomorphism property, that is,  $e$  is an edge in  $H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$  if and only if  $\psi(e)$  is an edge in  $(H_1 \widetilde{\times} H_2) \widetilde{\times} H_3$ . Let  $e = \{((x_1, y_1), z_1), \dots, ((x_r, y_r), z_r)\}$  be an edge in  $(H_1 \widetilde{\times} H_2) \widetilde{\times} H_3$ . There are two cases that can occur.

First,  $\{z_1, \dots, z_r\}$  is an edge in  $H_3$  and  $\{(x_1, y_1), \dots, (x_r, y_r)\}$  is therefore a subset of an edge in  $H_1 \widetilde{\times} H_2$ . Hence  $\{x_1, \dots, x_r\}$  and  $\{y_1, \dots, y_r\}$  must be subsets of edges in  $H_1$  and  $H_2$ , respectively. But then  $\{(y_1, z_1), \dots, (y_r, z_r)\}$  is an edge in  $H_2 \widetilde{\times} H_3$ , which implies that  $\psi(e) = \{(x_1, (y_1, z_1)), \dots, (x_r, (y_r, z_r))\}$  is an edge in  $H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$ .

Second,  $\{(x_1, y_1), \dots, (x_r, y_r)\}$  is an edge in  $H_1 \widetilde{\times} H_2$  and  $\{z_1, \dots, z_r\}$  is a subset of an edge in  $H_3$ . Then  $\{x_1, \dots, x_r\}$  is an edge in  $H_1$  and  $\{y_1, \dots, y_r\}$  is a subset of an edge in  $H_2$ , or vice versa. In the first case  $\{(y_1, z_1), \dots, (y_r, z_r)\}$  is a subset of an edge in  $H_2 \widetilde{\times} H_3$ , hence  $\psi(e)$  is an edge in  $H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$ , and in the second case  $\{(y_1, z_1), \dots, (y_r, z_r)\}$  is an edge in  $H_2 \widetilde{\times} H_3$  and thus  $\psi(e)$  is an edge in  $H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$ .

This implies that if  $e$  is an edge in  $(H_1 \widetilde{\times} H_2) \widetilde{\times} H_3$ , then  $\psi(e)$  is an edge in  $H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$ . The converse follows analogously. Thus  $(H_1 \widetilde{\times} H_2) \widetilde{\times} H_3 \cong H_1 \widetilde{\times} (H_2 \widetilde{\times} H_3)$ .  $\square$

**Proposition 3.** *The normal product  $\widetilde{\boxtimes}$  is associative.*

*Proof.* As in the previous proof, consider the bijection  $\psi : V((H_1 \widetilde{\boxtimes} H_2) \widetilde{\boxtimes} H_3) \rightarrow V(H_1 \widetilde{\boxtimes} (H_2 \widetilde{\boxtimes} H_3))$  defined as  $((x, y), z) \mapsto (x, (y, z))$ . We claim this is an isomorphism.

Let  $p_{1,2}$  be the projection from  $(H_1 \widetilde{\boxtimes} H_2) \widetilde{\boxtimes} H_3$  onto  $H_1 \widetilde{\boxtimes} H_2$ , defined by  $p_{1,2}(((x, y), z)) = (x, y)$ . Let  $p_{2,3}$  be projection from  $H_1 \widetilde{\boxtimes} (H_2 \widetilde{\boxtimes} H_3)$  to  $H_2 \widetilde{\boxtimes} H_3$ , whereas  $p_j$  is the usual projection to  $H_j$ . By definition,  $e = \{((x_1, y_1), z_1), \dots, ((x_r, y_r), z_r)\}$  is an edge in  $(H_1 \widetilde{\boxtimes} H_2) \widetilde{\boxtimes} H_3$  if and only if one of the following conditions is satisfied:

- (i)  $p_{1,2}(e) = e_{1,2} \in E(H_1 \widetilde{\boxtimes} H_2)$  and  $|p_3(e)| = 1$ ,
- (ii)  $p_3(e) = e_3 \in E(H_3)$  and  $|p_{1,2}(e)| = 1$ ,
- (iii)  $p_{1,2}(e) = e_{1,2} \in E(H_1 \widetilde{\boxtimes} H_2)$  and  $p_3(e) \subseteq e_3 \in E(H_3)$  and  $|e| = |e_{1,2}| = |p_{1,2}(e)| = |p_3(e)| \leq |e_3|$ ,
- (iv)  $p_3(e) = e_3 \in E(H_3)$  and  $p_{1,2}(e) \subseteq e_{1,2} \in E(H_1 \widetilde{\boxtimes} H_2)$  and  $|e| = |e_3| = |p_3(e)| = |p_{1,2}(e)| \leq |e_{1,2}|$ .

Condition (i) is equivalent to one of the following conditions holding:

- (i a)  $p_1(e) = p_1(e_{1,2}) = e_1 \in E(H_1)$  and  $|p_2(e)| = |p_2(e_{1,2})| = |p_3(e)| = 1$ , or
- (i b)  $p_2(e) = p_2(e_{1,2}) = e_2 \in E(H_2)$  and  $|p_1(e)| = |p_1(e_{1,2})| = |p_3(e)| = 1$ , or
- (i c)  $p_1(e) = p_1(e_{1,2}) = e_1 \in E(H_1)$  and  $p_2(e) = p_2(e_{1,2}) \subseteq e_2 \in E(H_2)$  and  $|e| = |e_1| = |p_1(e)| = |p_2(e)| \leq |e_2|$  and  $|p_3(e)| = 1$ , or
- (i d)  $p_2(e) = p_2(e_{1,2}) = e_2 \in E(H_2)$  and  $p_1(e) = p_1(e_{1,2}) \subseteq e_1 \in E(H_1)$  and  $|e| = |e_2| = |p_2(e)| = |p_1(e)| \leq |e_1|$  and  $|p_3(e)| = 1$ .

Condition (iii) is equivalent to one of the following conditions holding:

- (iii a)  $p_1(e) = p_1(e_{1,2}) = e_1 \in E(H_1)$  and  $|p_2(e)| = |p_2(e_{1,2})| = 1$  and  $p_3(e) \subseteq e_3 \in E(H_3)$  and  $|e| = |p_3(e)| \leq |e_3|$ , or
- (iii b)  $p_2(e) = p_2(e_{1,2}) = e_2 \in E(H_2)$  and  $|p_1(e)| = |p_1(e_{1,2})| = 1$  and  $p_3(e) \subseteq e_3 \in E(H_3)$  and  $|e| = |p_3(e)| \leq |e_3|$ , or
- (iii c)  $p_1(e) = p_1(e_{1,2}) = e_1 \in E(H_1)$  and  $p_2(e) = p_2(e_{1,2}) \subseteq e_2 \in E(H_2)$  and  $|e| = |e_1| = |p_1(e)| = |p_2(e)| \leq |e_2|$  and  $p_3(e) \subseteq e_3 \in E(H_3)$  and  $|e| = |p_3(e)| \leq |e_3|$ , or
- (iii d)  $p_2(e) = p_2(e_{1,2}) = e_2 \in E(H_2)$  and  $p_1(e) = p_1(e_{1,2}) \subseteq e_1 \in E(H_1)$  and  $|e| = |e_2| = |p_2(e)| = |p_1(e)| \leq |e_1|$  and  $p_3(e) \subseteq e_3 \in E(H_3)$  and  $|e| = |p_3(e)| \leq |e_3|$ .

Condition (iv) is equivalent to one of the following conditions holding:

- (iv a)  $p_3(e) = e_3 \in E(H_3)$  and  $p_1(e) = p_1(p_{1,2}(e)) \subseteq e_1 \in E(H_1)$  and  $|p_2(e)| = |p_2(p_{1,2}(e))| = 1$  and  $|e| = |p_3(e)| = |e_3| = |p_1(e)| \leq |e_1|$ , or
- (iv b)  $p_3(e) = e_3 \in E(H_3)$  and  $p_2(e) = p_2(p_{1,2}(e)) \subseteq e_2 \in E(H_2)$  and  $|p_1(e)| = |p_1(p_{1,2}(e))| = 1$  and  $|e| = |p_3(e)| = |e_3| = |p_2(e)| \leq |e_2|$ , or

- (iv c)  $p_3(e) = e_3 \in E(H_3)$  and  $p_1(e) = p_1(p_{1,2}(e)) \subseteq e_1 \in E(H_1)$  and  $p_2(e) = p_2(p_{1,2}(e)) \subseteq e_2 \in E(H_2)$  and  $|e| = |p_3(e)| = |e_3| = |p_1(e)| = |p_2(e)| \leq \min_{i=1,2} |e_i|$ .

Then Condition (i a) implies the following condition:

- (I)  $p_1(e) = e_1 \in E(H_1)$  and  $|p_{2,3}(e)| = 1$ .

Conditions (i b), (ii), (iii b) and (iv b) each imply the following condition:

- (II)  $|p_1(e)| = 1$  and  $p_{2,3}(e) = e_{2,3} \in E(H_2 \boxtimes H_3)$ .

Conditions (i c), (iii a) and (iii c) each imply the following condition:

- (III)  $p_1(e) = e_1 \in E(H_1)$  and  $p_{2,3}(e) \subseteq e_{2,3} \in E(H_2 \boxtimes H_3)$  and  $|e| = |e_1| = |p_1(e)| = |p_{2,3}(e)| \leq |e_{2,3}|$ .

Conditions (i d), (iii d), (iv a) and (iv c) each imply the following condition:

- (IV)  $p_1(e) \subseteq e_1 \in E(H_1)$  and  $p_{2,3}(e) = e_{2,3} \in E(H_2 \boxtimes H_3)$  and  $|e| = |e_{2,3}| = |p_{2,3}(e)| = |p_1(e)| \leq |e_1|$ .

By definition of the normal product, if any of the Conditions (I)–(IV) are satisfied, then  $\psi(e) = \{(x_1, (y_1, z_1)), \dots, (x_r, (y_r, z_r))\}$  is an edge in  $H_1 \boxtimes (H_2 \boxtimes H_3)$ . It follows that  $\psi$  is a homomorphism. In the same way, the inverse  $(x, (y, z)) \mapsto ((x, y), z)$  is also a homomorphism.  $\square$

## References

- [1] C. Berge. *Hypergraphs: Combinatorics of finite sets*, volume 45. North-Holland, Amsterdam, 1989.
- [2] Timothy Black. Monotone properties of  $k$ -uniform hypergraphs are weakly evasive. In *Proceedings of the 2015 Conference on Innovations in Theoretical Computer Science, ITCS '15*, pages 383–391, New York, NY, USA, 2015. ACM.
- [3] A. Bretto, Y. Silvestre, and T. Vallée. Cartesian product of hypergraphs: properties and algorithms. In *4th Athens Colloquium on Algorithms and Complexity (ACAC 2009)*, volume 4 of *EPTCS*, pages 22–28, 2009.
- [4] W. Dörfler. Multiple Covers of Hypergraphs. *Annals of the New York Academy of Sciences*, 319(1):169–176, 1979.
- [5] R. Hammack, W. Imrich, and S. Klavžar. *Handbook of Product Graphs*. Discrete Mathematics and its Applications. CRC Press, 2nd edition, 2011.
- [6] M. Hellmuth and F. Lehner. Fast factorization of cartesian products of (directed) hypergraphs. *J. Theor. Comp. Sci.*, 2015. submitted.
- [7] M. Hellmuth, L. Ostermeier, and M. Noll. Strong products of hypergraphs: Unique prime factorization theorems and algorithms. *Discr. Appl. Math.*, 171:60–71, 2014.
- [8] M. Hellmuth, L. Ostermeier, and P. F. Stadler. A survey on hypergraph products. *Math. Comp. Sci.*, 6:1–32, 2012.
- [9] W. Imrich. Kartesisches Produkt von Mengensystemen und Graphen. *Studia Sci. Math. Hungar.*, 2:285 – 290, 1967.
- [10] W. Imrich. über das schwache Kartesische Produkt von Graphen. *Journal of Combinatorial Theory*, 11(1):1–16, 1971.
- [11] W. Imrich and H. Izicki. Associative products of graphs. *Monatshefte für Mathematik*, 80(4):277–281, 1975.
- [12] A. Kaveh and B. Alinejad. Hypergraph products for structural mechanics. In B.H.V. Topping, editor, *Proceedings of the Eleventh International Conference on Computational Structures Technology*, Stirlingshire, UK, 2012. Civil-Comp Press. Paper 266.
- [13] A. Kaveh and B. Alinejad. Hypergraph products for structural mechanics. *Advances in Engineering Software*, 80:72 – 81, 2015. Civil-Comp.
- [14] L. Ostermeier, M. Hellmuth, and P. F. Stadler. The Cartesian product of hypergraphs. *Journal of Graph Theory*, 2011.

- [15] M. Sonntag. Hamiltonian properties of the Cartesian sum of hypergraphs. *J. Inf. Process. Cybern.*, 25(3):87–100, 1989.
- [16] M. Sonntag. Hamiltonicity of the normal product of hypergraphs. *J. Inf. Process. Cybern.*, 26(7):415–433, 1990.
- [17] M. Sonntag. *Hamiltonsche Eigenschaften von Produkten von Hypergraphen*. Habilitation, Fakultät für Mathematik und Naturwissenschaften, Bergakademie Freiberg, 1991.
- [18] X. Zhu. On the chromatic number of the product of hypergraphs. *Ars Comb.*, 34:25–31, 1992.

Richard H. Hammack  
 Department of Mathematics and Applied Mathematics  
 Virginia Commonwealth University  
 Richmond, VA 23284-2014, USA  
 e-mail: [rhammack@vcu.edu](mailto:rhammack@vcu.edu)

Marc Hellmuth  
 Department of Mathematics and Computer Science  
 University of Greifswald  
 Walther-Rathenau-Straße 47, D-17487 Greifswald, Germany  
 Center for Bioinformatics  
 Saarland University  
 Building E 2.1, Room 413, P.O. Box 15 11 50, D-66041 Saarbrücken, Germany  
 e-mail: [mhellmuth@mailbox.org](mailto:mhellmuth@mailbox.org)

Lydia Ostermeier  
 Bioinformatics Group,  
 Department of Computer Science and Interdisciplinary Center for Bioinformatics  
 University of Leipzig,  
 Härtelstraße 16-18, D-04107 Leipzig, Germany  
 e-mail: [glydia@bioinf.uni-leipzig.de](mailto:glydia@bioinf.uni-leipzig.de)

Peter F. Stadler  
 Bioinformatics Group,  
 Department of Computer Science; and Interdisciplinary Center for Bioinformatics,  
 University of Leipzig,  
 Härtelstraße 16-18, D-04107 Leipzig, Germany  
 Max Planck Institute for Mathematics in the Sciences  
 Inselstrasse 22, D-04103 Leipzig, Germany  
 RNomics Group, Fraunhofer Institut für Zelltherapie und Immunologie, Deutscher Platz 5e, D-04103 Leipzig, Germany  
 Department of Theoretical Chemistry, University of Vienna, Währingerstraße 17, A-1090 Wien, Austria  
 Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM87501, USA  
 e-mail: [studla@bioinf.uni-leipzig.de](mailto:studla@bioinf.uni-leipzig.de)